

Deep Learning

Tutorial 1 – Vector Calculus

Naama Pearl
Autonomous Vision Group
University of Tübingen

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Tübingen AI Center
tuebingen.ai



e l i a s
European Laboratory for Learning and Intelligent Systems

Outline

- ▶ Quiz
- ▶ Organization
- ▶ Derivatives in 1D
- ▶ Partial derivatives and gradients
- ▶ Jacobians
- ▶ Working problems

Quiz



Tutorials and Exercise Sheets

Tutorials

- ▶ Every **Wednesday, 16:15–17:45**, same room as the lecture (after a 30 min break)
- ▶ Tutors: Yong Cao, Glenn Angrabeit, Naama Pearl
- ▶ Hands-on and interactive:
 - ▶ A short **quiz** at the beginning
 - ▶ Background you need for the sheets (math, Python, NumPy, ...)
 - ▶ **Working problems**: you solve, then we go through the solutions together
 - ▶ Introducing a new sheet, or discussing the solutions of the last one
- ▶ **Ask questions!** In the tutorial, in the Ilias forum, or at the weekly helpdesk

Exercise sheets

- ▶ **6 sheets**, one every **two weeks**, on the [course website](#)
- ▶ Each sheet has **pen & paper** problems and **coding** problems (Python notebooks)
- ▶ You solve them **at home**
- ▶ **Nothing is handed in.** We publish the solutions and discuss them in the tutorial
- ▶ Solving the sheets on your own is good **preparation for the exam**

Date	Tutorial	Sheet
14.10. (today)	1: Vector calculus	Sheet 1 out
21.10.	2: Python, NumPy and coding the sheet	work on sheet 1
28.10.	3	Solutions 1 , sheet 2 out

Exercise sheet 1

[Sheet 1 \(PDF\)](#)

Exercise	You need	Covered in
<i>Pen & paper</i>		
1. Derivatives	derivatives, Jacobians, chain rule	tutorial 1 (today)
2. Linear regression	linear regression, partial derivatives	lecture 1, tutorial 1 (today)
3. Computation graphs	computation graphs, backpropagation	lecture 2 (next week)
4. Logistic regression	logistic regression, chain rule	lecture 1 or 2
<i>Coding</i>		
1. Vector derivative	NumPy, broadcasting	tutorial 2 (next week)
2. Linear regression	NumPy, the coding framework	lecture 2, tutorial 2 (next week)
3. Logistic regression	NumPy, the coding framework	lecture 2, tutorial 2 (next week)

- Ask in the **forum** or come to the **helpdesk**
- **Solutions** are published and discussed in the tutorial on **28.10.**

Next week

Bring your laptop

Fully charged. Tutorial 2 is a coding tutorial

Setup for tutorial 2

- ▶ Tutorial 2 is a **coding** tutorial: Python, NumPy and the coding part of sheet 1
- ▶ **Bring your laptop** (charged)
- ▶ **Before next week**, set up the Python environment for sheet 1:

1. Install `conda`:

`https://docs.conda.io/projects/conda/en/latest/user-guide/install/`

2. In the sheet's code folder: `conda env create -f environment.yml`

3. `conda activate deep_learning_ex_1` and then `jupyter lab`

If step 2 is stuck at *Solving environment*, try adding `--solver libmamba` (it can still take a few minutes)

- ▶ Problems with the setup? Come to the helpdesk or ask in the forum **before** the tutorial

Derivatives in 1D

What is a derivative?


- The **derivative** of $f: \mathbb{R} \rightarrow \mathbb{R}$ at a point x measures how sensitive the output is to a small change of the input:

$$\frac{\text{change of output}}{\text{change of input}} \longrightarrow f'(x) = \frac{df}{dx}(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- In deep learning: how much does the **loss** change if we change a **weight** a little?
 $d\mathcal{L}/dw$

The derivative is the slope

We can visualize the derivative as the slope of the function at a point. For the loss $\mathcal{L}$ at the weight w , the slope is $\frac{d\mathcal{L}}{dw}$.



- Here the slope is **positive**: the loss grows with w
- So we push the weight to the **left** (make w smaller) to decrease the loss
- Gradient descent:
$$w_{\text{new}} = w - \eta \frac{d\mathcal{L}}{dw}$$
- With a negative slope the step goes to the right

Shape of the derivative

Before computing the derivative:

1. What is the **output** dimension of the function?
2. What is the **input** dimension of the variable we differentiate with respect to?
3. What is the **shape of the derivative**?

Throughout the tutorial: **output dimensions in red**, **input dimensions in blue**.

Dimensions in 1D

- ▶ $f: \mathbb{R}^1 \rightarrow \mathbb{R}^1$: scalar input, scalar output
- ▶ The derivative is a scalar

$$f: \mathbb{R}^1 \rightarrow \mathbb{R}^1 \quad x \in \mathbb{R}^1 \quad \Rightarrow \quad f'(x) = \frac{df}{dx}(x) \in \mathbb{R}^1$$

Example: derivative from the definition

$$f(x) = x^2, \quad f: \mathbb{R}^1 \rightarrow \mathbb{R}^1 \quad \Rightarrow \quad f'(x) \in \mathbb{R}^1$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h}$$

plug in f

$$= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h}$$

expand

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h}$$

$$x^2 - x^2 = 0$$

$$= \lim_{h \rightarrow 0} (2x + h)$$

divide by h

$$= 2x$$

let $h \rightarrow 0$

Derivative rules

Elementary functions

$f(x)$	$f'(x)$
c	0
x^n	$n x^{n-1}$
e^x	e^x
$\log x$	$\frac{1}{x}$
$\sin x$	$\cos x$
$\cos x$	$-\sin x$

Combinations

Sum: $(u + v)' = u' + v'$

Constant factor: $(c u)' = c u'$

Product: $(u v)' = u' v + u v'$


Quotient: $\left(\frac{u}{v}\right)' = \frac{u' v - u v'}{v^2}$

Chain: $(f(g(x)))' = f'(g(x)) g'(x)$

The **chain rule** is the core of how backpropagation computes gradients.

Derivative of a neuron (chain rule)

$$f(w) = \tanh(wx + b), \quad x, b \text{ fixed}, \quad f: \mathbb{R}^1 \rightarrow \mathbb{R}^1 \quad \Rightarrow \quad \frac{df}{dw} \in \mathbb{R}^1$$



$$f(w) = \tanh(wx + b)$$

$$= g(u(w)) \quad \text{with} \quad g(u) = \tanh u, \quad u(w) = wx + b \quad \text{activation of a linear layer}$$

$$\frac{df}{dw} = g'(u) u'(w)$$

chain rule

$$= (1 - \tanh^2(wx + b)) \cdot x$$

$$\tanh'(u) = 1 - \tanh^2 u, \quad u'(w) = x$$

- In backpropagation, $d\mathcal{L}/df$ comes from the layers above, and we multiply by the **local** derivatives: $\frac{d\mathcal{L}}{dw} = \frac{d\mathcal{L}}{df} \frac{df}{du} \frac{du}{dw}$


Derivative of a ReLU neuron

Exercise

Same neuron, now with a **ReLU** activation (x, b fixed):

$$f(w) = \text{ReLU}(w x + b)$$

$$\text{ReLU}(u) = \max(0, u) = \begin{cases} 0 & u < 0 \\ u & u \geq 0 \end{cases}$$



1. Compute $\frac{df}{dw}$.

Reminder (chain rule): $f = g(u(w)) \Rightarrow \frac{df}{dw} = g'(u) u'(w)$



Solution: derivative of a ReLU neuron

$$f(w) = \max(0, w x + b), \quad x, b \text{ fixed}, \quad f: \mathbb{R}^1 \rightarrow \mathbb{R}^1 \quad \Rightarrow \quad \frac{df}{dw} \in \mathbb{R}^1$$

$$f(w) = g(u(w)), \quad g(u) = \max(0, u), \quad u(w) = w x + b$$

$$\frac{df}{dw} = g'(u) u'(w) = g'(u) \cdot x$$

$$= \begin{cases} 1 \cdot x = x & w x + b > 0 \\ 0 \cdot x = 0 & w x + b < 0 \end{cases}$$



- ▶ Same chain rule as for $\tanh$, only g' changes
- ▶ At $w x + b = 0$: not defined
(in practice, frameworks use 0)

Partial Derivatives and Gradients

Partial derivatives of a neuron

$$f(w, b) = \tanh(w x + b), \quad x \text{ fixed}, \quad f: \mathbb{R}^2 \rightarrow \mathbb{R}^1 \quad \Rightarrow \quad \nabla_{(w,b)} f \in \mathbb{R}^2$$

$$\frac{\partial f}{\partial w} = \frac{\partial}{\partial w} \tanh(w x + b) = (1 - \tanh^2(w x + b)) \cdot x \quad \text{as before: } b \text{ is a constant}$$

$$\frac{\partial f}{\partial b} = \frac{\partial}{\partial b} \tanh(w x + b) = (1 - \tanh^2(w x + b)) \cdot 1 \quad \text{now } w \text{ is a constant}$$

$$\nabla_{(w,b)} f = \begin{pmatrix} \frac{\partial f}{\partial w} \\ \frac{\partial f}{\partial b} \end{pmatrix} = (1 - \tanh^2(w x + b)) \begin{pmatrix} x \\ 1 \end{pmatrix} \in \mathbb{R}^2 \quad \text{stack them in a column}$$

- When we computed the derivative w.r.t. the weight w before, we kept b fixed. This was a **partial derivative** (written ∂ instead of d)
- The **gradient** has the shape of the parameters (w, b)

Partial derivatives and the gradient

- ▶ Let $f: \mathbb{R}^n \rightarrow \mathbb{R}^1$ depend on several inputs $x_1, \dots, x_n$
- ▶ The **partial derivative** $\frac{\partial f}{\partial x_j}$: differentiate w.r.t. x_j and treat all other variables as constants
- ▶ The **gradient** collects all n partial derivatives in a column, in the **shape of $\mathbf{x}$** :

$$\nabla_{\mathbf{x}} f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f}{\partial x_1}(\mathbf{x}) \\ \vdots \\ \frac{\partial f}{\partial x_n}(\mathbf{x}) \end{pmatrix} \in \mathbb{R}^n$$

$$f: \mathbb{R}^n \rightarrow \mathbb{R}^1 \quad \mathbf{x} \in \mathbb{R}^n \quad \Rightarrow \quad \nabla_{\mathbf{x}} f \in \mathbb{R}^n \text{ (same shape as } \mathbf{x} \text{)}$$

- ▶ In deep learning: the gradient of the loss w.r.t. the weights has the shape of the weights, so we can use it to update them: $\mathbf{w}^{t+1} = \mathbf{w}^t - \eta \nabla_{\mathbf{w}} \mathcal{L}$

Partial derivatives and gradient

Exercise

A neuron with two inputs ($\mathbf{x}$ fixed):

$$\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad f(\mathbf{w}) = \tanh(\mathbf{w}^\top \mathbf{x})$$

1. Shapes: $f: \mathbb{R}^? \rightarrow \mathbb{R}^?$, $\nabla_{\mathbf{w}} f \in ?$
2. Compute $\frac{\partial f}{\partial w_1}$, $\frac{\partial f}{\partial w_2}$ and $\nabla_{\mathbf{w}} f$.

Reminder: $\tanh'(u) = 1 - \tanh^2 u$

Solution: $f(\mathbf{w}) = \tanh(\mathbf{w}^\top \mathbf{x})$

$$\mathbf{w}, \mathbf{x} \in \mathbb{R}^2, \quad f: \mathbb{R}^? \rightarrow \mathbb{R}^? \quad \Rightarrow \quad \nabla_{\mathbf{w}} f \in ? \quad \text{What are the shapes?}$$

$$f(\mathbf{w}) = \tanh(w_1 x_1 + w_2 x_2)$$

$\mathbf{w}^\top \mathbf{x}$ in components

$$\frac{\partial f}{\partial w_1} = (1 - \tanh^2(\mathbf{w}^\top \mathbf{x})) \cdot x_1$$

chain rule, w_2 constant

$$\frac{\partial f}{\partial w_2} = (1 - \tanh^2(\mathbf{w}^\top \mathbf{x})) \cdot x_2$$

chain rule, w_1 constant

$$\nabla_{\mathbf{w}} f = (1 - \tanh^2(\mathbf{w}^\top \mathbf{x})) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = (1 - \tanh^2(\mathbf{w}^\top \mathbf{x})) \mathbf{x} \in \mathbb{R}^2 \quad \text{stack in a column, shape of } \mathbf{w} \checkmark$$

The gradient w.r.t. the weights is the input $\mathbf{x}$, times a scalar.

Gradients of Vector Expressions

How to differentiate vector expressions

For $\tanh(\mathbf{w}^\top \mathbf{x})$ we wrote $\mathbf{w}^\top \mathbf{x}$ in components. This works for any scalar expression of a vector, like $\mathbf{w}^\top \mathbf{x}$, $\mathbf{w}^\top \mathbf{w}$, $\mathbf{x}^\top \mathbf{A} \mathbf{x}$, ...

1. **Shapes:** f is a scalar, $\mathbf{x} \in \mathbb{R}^n \Rightarrow \nabla_{\mathbf{x}} f \in \mathbb{R}^n$ (shape of $\mathbf{x}$)
2. **Components:** write f as a sum of scalars, e.g. $\mathbf{w}^\top \mathbf{x} = \sum_i w_i x_i$
3. **Scalar derivative:** compute $\frac{\partial f}{\partial x_j}$ with the 1D rules (everything else is a constant)
4. **Reassemble:** put the n entries into a column and check the shape

Sanity check: for $n = 1$ the result must reduce to the 1D derivative.

Example: $f(\mathbf{x}) = \mathbf{w}^\top \mathbf{x}$

$$f(\mathbf{x}) = \mathbf{w}^\top \mathbf{x}, \quad \mathbf{w}, \mathbf{x} \in \mathbb{R}^n, \quad f: \mathbb{R}^? \rightarrow \mathbb{R}^? \quad \Rightarrow \quad \nabla_{\mathbf{x}} f \in ? \quad \text{What are the shapes?}$$

$$f(\mathbf{x}) = \sum_{i=1}^n w_i x_i$$

components

$$\frac{\partial f}{\partial x_j} = \sum_{i=1}^n w_i \frac{\partial x_i}{\partial x_j}$$

scalar derivative w.r.t. one x_j

$$= \sum_{i=1}^n w_i \cdot \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} = w_j$$

only the term $i = j$ survives

$$\nabla_{\mathbf{x}} f = \begin{pmatrix} w_1 & \cdots & w_n \end{pmatrix}^\top = \mathbf{w} \in \mathbb{R}^n$$

reassemble as a column, shape $\checkmark$

Sanity check $n = 1$: $f(x) = wx \Rightarrow f'(x) = w \checkmark$

Gradients of $\mathbf{w}^\top \mathbf{x}$ and $\mathbf{w}^\top \mathbf{w}$

Exercise

Let $\mathbf{w}, \mathbf{x} \in \mathbb{R}^n$.

1. The same expression as a function of $\mathbf{w}$, $f(\mathbf{w}) = \mathbf{w}^\top \mathbf{x}$:
compute $\nabla_{\mathbf{w}} f$
2. $g(\mathbf{w}) = \mathbf{w}^\top \mathbf{w} = \|\mathbf{w}\|_2^2$:
compute $\nabla_{\mathbf{w}} g$

Previous example: $f(\mathbf{x}) = \mathbf{w}^\top \mathbf{x}$

$$f(\mathbf{x}) = \sum_{i=1}^n w_i x_i$$

$$\frac{\partial f}{\partial x_j} = \sum_{i=1}^n w_i \frac{\partial x_i}{\partial x_j} = \sum_{i=1}^n w_i \cdot \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} = w_j$$

$$\nabla_{\mathbf{x}} f = \begin{pmatrix} w_1 & \cdots & w_n \end{pmatrix}^\top = \mathbf{w}$$

Solution: gradients of $\mathbf{w}^\top \mathbf{x}$ and $\mathbf{w}^\top \mathbf{w}$

$\mathbf{w}, \mathbf{x} \in \mathbb{R}^n$, $f, g \in \mathbb{R}^?$ $\Rightarrow \nabla_{\mathbf{w}} f, \nabla_{\mathbf{w}} g \in ?$ What are the shapes?

$$f = \sum_{i=1}^n w_i x_i$$

components

$$\frac{\partial f}{\partial w_j} = \sum_{i=1}^n \frac{\partial w_i}{\partial w_j} x_i = \sum_{i=1}^n \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \cdot x_i = x_j$$

now $\mathbf{x}$ is the constant

$$\nabla_{\mathbf{w}} f = \begin{pmatrix} x_1 & \cdots & x_n \end{pmatrix}^\top = \mathbf{x}$$

(before: $\nabla_{\mathbf{x}} f = \mathbf{w}$)

$$g = \sum_{i=1}^n w_i w_i = \sum_{i=1}^n w_i^2$$

components

$$\frac{\partial g}{\partial w_j} = \sum_{i=1}^n 2w_i \frac{\partial w_i}{\partial w_j} = \sum_{i=1}^n 2w_i \cdot \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} = 2w_j$$

chain rule for w_i^2 , only the term $i = j$ survives

$$\nabla_{\mathbf{w}} g = \begin{pmatrix} 2w_1 & \cdots & 2w_n \end{pmatrix}^\top = 2\mathbf{w}$$

Jacobians and the Chain Rule in Matrix Form

A function with several outputs

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad \mathbf{f}(\mathbf{x}) = \begin{pmatrix} x_1 x_2 \\ x_1 + x_2^2 \\ e^{x_1} \end{pmatrix}, \quad \mathbf{f}: \mathbb{R}^? \rightarrow \mathbb{R}^? \quad \Rightarrow \quad \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \in ?$$

What are the shapes?

$$\frac{\partial \mathbf{f}}{\partial \mathbf{x}} = \begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} \\ \frac{\partial f_3}{\partial x_1} & \frac{\partial f_3}{\partial x_2} \end{pmatrix} = \begin{pmatrix} x_2 & x_1 \\ 1 & 2x_2 \\ e^{x_1} & 0 \end{pmatrix}$$

- Each output gets one row with its partial derivatives: row 1 for $f_1 = x_1 x_2$, row 2 for f_2 , row 3 for f_3
- Stacking the rows gives a 3×2 matrix (3 outputs, 2 inputs): the **Jacobian**

Derivative of a vector-to-vector function: the Jacobian

In general, let $\mathbf{f}: \mathbb{R}^n \rightarrow \mathbb{R}^m$, i.e., n inputs and m outputs $\mathbf{f}(\mathbf{x}) = (f_1(\mathbf{x}), \dots, f_m(\mathbf{x}))^\top$.

$$\frac{\partial \mathbf{f}}{\partial \mathbf{x}}(\mathbf{x}) = \mathbf{J}(\mathbf{x}) = \begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \dots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \dots & \frac{\partial f_m}{\partial x_n} \end{pmatrix} \in \mathbb{R}^{m \times n}$$

- ▶ Row i belongs to output f_i : all partial derivatives of $f_i \Rightarrow m$ rows
- ▶ Column j belongs to input x_j : how all outputs react to $x_j \Rightarrow n$ columns
- ▶ Entry (i, j) is an ordinary scalar derivative: $\frac{\partial f_i}{\partial x_j}$

Gradient and Jacobian

Jacobian $\frac{\partial \mathbf{f}}{\partial \mathbf{x}}$

- ▶ “If I change the input a little, how much does the **output** change?”
- ▶ $d\mathbf{f} \approx \frac{\partial \mathbf{f}}{\partial \mathbf{x}} d\mathbf{x}$
- ▶ Shape: **#outputs** $\times$ **#inputs**
(for a scalar f : a row)
- ▶ Used in the **chain rule**

Gradient $\nabla_{\mathbf{x}} f$ (scalar f only)

- ▶ All partial derivatives $\frac{\partial f}{\partial x_j}$,
arranged in the **shape of $\mathbf{x}$**
- ▶ So it is easy to update $\mathbf{x}$:
 $\mathbf{x} \leftarrow \mathbf{x} - \eta \nabla_{\mathbf{x}} f$
- ▶ Used in **gradient descent**

For a scalar f , they hold the same numbers, so $\nabla_{\mathbf{x}} f = \left(\frac{\partial f}{\partial \mathbf{x}} \right)^\top$.

Example: $f(\mathbf{x}) = \mathbf{w}^\top \mathbf{x}$: $\frac{\partial f}{\partial \mathbf{x}} = \mathbf{w}^\top$ (row), $\nabla_{\mathbf{x}} f = \mathbf{w}$ (column)

Shapes of derivatives

function	w.r.t.	derivative	name
$f \in \mathbb{R}^1$	$x \in \mathbb{R}^1$	$\frac{\partial f}{\partial x} \in \mathbb{R}^1$	derivative
$f \in \mathbb{R}^1$	$\mathbf{x} \in \mathbb{R}^n$	$\nabla_{\mathbf{x}} f \in \mathbb{R}^n$	gradient (shape of $\mathbf{x}$)
$\mathcal{L} \in \mathbb{R}^1$	$\mathbf{W} \in \mathbb{R}^{m \times n}$	$\nabla_{\mathbf{W}} \mathcal{L} \in \mathbb{R}^{m \times n}$	gradient (shape of $\mathbf{W}$)
$\mathbf{f} \in \mathbb{R}^m$	$\mathbf{x} \in \mathbb{R}^n$	$\frac{\partial \mathbf{f}}{\partial \mathbf{x}} \in \mathbb{R}^{m \times n}$	Jacobian

- There is **one number for every pair (output, input)**

Shape and first row of a Jacobian

Exercise

$$\mathbf{f}(\mathbf{x}) = \begin{pmatrix} x_1 x_2 x_3 \\ x_1 + x_3 \end{pmatrix}$$

1. What is the shape of $\frac{\partial \mathbf{f}}{\partial \mathbf{x}}$?
2. Compute its first row.

Solution: Jacobian of $\mathbf{f}$

$$\mathbf{f}(\mathbf{x}) = \begin{pmatrix} x_1 x_2 x_3 \\ x_1 + x_3 \end{pmatrix}, \quad \mathbf{f}: \mathbb{R}^? \rightarrow \mathbb{R}^? \quad \Rightarrow \quad \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \in ? \quad \text{What is the shape?}$$

$$\frac{\partial \mathbf{f}}{\partial \mathbf{x}} = \begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \frac{\partial f_1}{\partial x_3} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \frac{\partial f_2}{\partial x_3} \end{pmatrix} = \begin{pmatrix} x_2 x_3 & x_1 x_3 & x_1 x_2 \\ 1 & 0 & 1 \end{pmatrix}$$

- ▶ 2 outputs (rows), 3 inputs (columns) $\Rightarrow 2 \times 3$
- ▶ Row 1: partial derivatives of $f_1 = x_1 x_2 x_3$ (the other two variables are constants)
- ▶ Row 2: partial derivatives of $f_2 = x_1 + x_3$ (it does not depend on x_2)

Example: least squares with the chain rule

$$\mathcal{L}(\mathbf{w}) = \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2, \quad \mathbf{X} \in \mathbb{R}^{N \times n}, \quad \mathbf{w} \in \mathbb{R}^n, \quad \mathbf{y} \in \mathbb{R}^N, \quad \mathcal{L}: \mathbb{R}^? \rightarrow \mathbb{R}^? \Rightarrow \nabla_{\mathbf{w}} \mathcal{L} \in ?$$

What are the shapes?

$$\mathbf{u} = \mathbf{X}\mathbf{w} - \mathbf{y} \in \mathbb{R}^N, \quad \mathcal{L} = \mathbf{u}^\top \mathbf{u}$$

inner (vector), outer (scalar)

$$\frac{\partial \mathcal{L}}{\partial \mathbf{w}} = \frac{\partial \mathcal{L}}{\partial \mathbf{u}} \frac{\partial \mathbf{u}}{\partial \mathbf{w}} = \underbrace{2\mathbf{u}^\top}_{1 \times N} \underbrace{\mathbf{X}}_{N \times n} \in \mathbb{R}^{1 \times n}$$

chain rule (details on the next slide)

$$\nabla_{\mathbf{w}} \mathcal{L} = \left(\frac{\partial \mathcal{L}}{\partial \mathbf{w}} \right)^\top = \mathbf{X}^\top 2(\mathbf{X}\mathbf{w} - \mathbf{y}) \in \mathbb{R}^n$$

the gradient is the row as a column

Same as in lecture 1: $\nabla_{\mathbf{w}} \mathcal{L} = 2\mathbf{X}^\top \mathbf{X}\mathbf{w} - 2\mathbf{X}^\top \mathbf{y}$ ✓

Least squares: details

$$\mathcal{L}(\mathbf{w}) = \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2, \quad \mathbf{X} \in \mathbb{R}^{N \times n}, \quad \mathbf{w} \in \mathbb{R}^n, \quad \mathbf{y} \in \mathbb{R}^N, \quad \mathcal{L}: \mathbb{R}^n \rightarrow \mathbb{R}^1 \Rightarrow \nabla_{\mathbf{w}} \mathcal{L} \in \mathbb{R}^n$$

$$\mathbf{u} = \mathbf{X}\mathbf{w} - \mathbf{y} \in \mathbb{R}^N, \quad \mathcal{L} = \mathbf{u}^\top \mathbf{u}$$

inner (vector), outer (scalar)

$$\frac{\partial \mathcal{L}}{\partial \mathbf{u}} = 2\mathbf{u}^\top \in \mathbb{R}^{1 \times N}$$

as for $\mathbf{w}^\top \mathbf{w}$, written as a row (1 output, N inputs)

$$\frac{\partial \mathbf{u}}{\partial \mathbf{w}} = \frac{\partial}{\partial \mathbf{w}} \begin{pmatrix} \mathbf{x}_1^\top \mathbf{w} - y_1 \\ \vdots \\ \mathbf{x}_N^\top \mathbf{w} - y_N \end{pmatrix} = \begin{pmatrix} \mathbf{x}_1^\top \\ \vdots \\ \mathbf{x}_N^\top \end{pmatrix} = \mathbf{X} \in \mathbb{R}^{N \times n}$$

row i : one sample, as for $\mathbf{w}^\top \mathbf{x}$

$$\frac{\partial \mathcal{L}}{\partial \mathbf{w}} = \frac{\partial \mathcal{L}}{\partial \mathbf{u}} \frac{\partial \mathbf{u}}{\partial \mathbf{w}} = \underbrace{2\mathbf{u}^\top}_{1 \times N} \underbrace{\mathbf{X}}_{N \times n} \in \mathbb{R}^{1 \times n}$$

chain rule: multiply, the inner N matches

$$\nabla_{\mathbf{w}} \mathcal{L} = \left(\frac{\partial \mathcal{L}}{\partial \mathbf{w}} \right)^\top = \mathbf{X}^\top 2(\mathbf{X}\mathbf{w} - \mathbf{y}) \in \mathbb{R}^n$$

the gradient is this row as a column (shape of $\mathbf{w}$)

Chain rule for vector-to-vector functions

In general, let $\mathbf{f}: \mathbb{R}^n \rightarrow \mathbb{R}^m$, $\mathbf{g}: \mathbb{R}^m \rightarrow \mathbb{R}^p$ and $\mathbf{h}(\mathbf{x}) = \mathbf{g}(\mathbf{f}(\mathbf{x}))$, so $\mathbf{h}: \mathbb{R}^n \rightarrow \mathbb{R}^p$. Then

$$\underbrace{\frac{\partial \mathbf{h}}{\partial \mathbf{x}}(\mathbf{x})}_{p \times n} = \underbrace{\frac{\partial \mathbf{g}}{\partial \mathbf{y}}(\mathbf{f}(\mathbf{x}))}_{p \times m} \underbrace{\frac{\partial \mathbf{f}}{\partial \mathbf{x}}(\mathbf{x})}_{m \times n}$$

- ▶ The 1D chain rule, but now a **matrix product**
- ▶ The inner dimension m (output of $\mathbf{f}$ = input of $\mathbf{g}$) always matches
- ▶ The result has the **output** dimension of $\mathbf{g}$ and the **input** dimension of $\mathbf{f}$
- ▶ Scalar loss ($p = 1$) with gradients, as for least squares: $\underbrace{\nabla_{\mathbf{x}} h}_{n \times 1} = \underbrace{\left(\frac{\partial \mathbf{f}}{\partial \mathbf{x}} \right)^{\top}}_{n \times m} \underbrace{\nabla_{\mathbf{y}} g}_{m \times 1}$, i.e.

$$\frac{\partial h}{\partial x_j} = \sum_{i=1}^m \frac{\partial g}{\partial y_i} \frac{\partial f_i}{\partial x_j} \quad (\text{multivariate chain rule, lecture 2})$$

Next week

Bring your laptop

Fully charged. Tutorial 2 is a coding tutorial

Summary

- ▶ Derivative = how much the output changes for a small change of the input. Its sign tells gradient descent which way to move
- ▶ Partial derivative: differentiate w.r.t. one variable, all others are constants
- ▶ Scalar function: the gradient $\nabla_{\mathbf{x}} f \in \mathbb{R}^n$ is in the shape of $\mathbf{x}$
- ▶ Differentiating vector expressions: shapes $\rightarrow$ components $\rightarrow$ scalar derivative $\rightarrow$ reassemble
- ▶ Shape of a Jacobian:

$$\mathbf{f}: \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \Rightarrow \quad \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \in \mathbb{R}^{m \times n} \quad (\text{rows} = \text{outputs}, \text{columns} = \text{inputs})$$

- ▶ Chain rule = product of Jacobians, the inner dimensions match:
 $(p \times n) = (p \times m)(m \times n)$. Scalar output: $\nabla_{\mathbf{x}} h = \left(\frac{\partial \mathbf{f}}{\partial \mathbf{x}}\right)^\top \nabla_{\mathbf{y}} g$